



Variability of Long Term Cosmic Ray: A Study of Solar Cycle 22 To 24

Sangita Mahendra Rajput, Research Scholar, Department of Electronics Engineering, SunRise University Alwar, Rajasthan
Dr. Ram Mohan Singh Bhadoriya, Associate Professor, Department of Electronics Engineering, SunRise University Alwar, Rajasthan

Abstract

In this paper we have studied the cosmic ray variability during solar cycle 22 to 24. For our study we have taken yearly average value of sunspot number (Rz) and cosmic ray intensity of two different latitude neutron monitor. We observed that trend of solar activity during period of solar cycle 22 to 24 have decreased and their consequence as well as the cosmic ray intensity have increased during this time period. We found that solar cycle 24 is the weakest solar cycle than previously occurred two solar cycles i.e. 22 and 23.

Key words - Sunspot Number (Rz). Cosmic Ray Intensity.

Introduction - Study of cosmic ray modulation with solar activity parameters is old and so on. Cosmic ray variation is based on the solar activity parameters. It has been known for a long time that intensity as well as the energy spectrum of galactic cosmic ray is modulated by solar activity. Among the various solar activity parameters, the sunspot number has been considered as a primary indicator to define the level of solar activity which generally follows 11-years periodicity. The sunspot numbers are used as solar index for study of cosmic ray modulation and solar terrestrial relationship (Dorman & Dorman 1967; Pomerantz & Duggal 1971; Rao 1972; Webber and Lockwood 1988; Ahluwalia 1998). Similarly several solar indices (solar flares, solar flux, coronal index, etc.) have been used as proxy index to represent the solar activity (Kane 2005; Gupta *et al.* 2005, 2006). Cosmic ray is anticorrelated with solar activity parameters (Ahluwalia & Wilson 1996; Dorman *et al.* 2001; Usoskin *et al.* 1998). Galactic cosmic ray in the energy range from several hundred MeV to tens of GeV is subjected to heliospheric modulation, under the influence of solar output and its variation. The heliospheric modulation of cosmic ray intensity and spectrum are associated with 11-year solar activity cycle. The charge/ polarity dependence of drift mechanism is clearly observed in cosmic ray modulation in terms of 22-year solar magnetic cycle, showing different shape of cosmic ray maxima in the alternate solar cycles. Long-term cosmic ray modulation in the high energy range is studied using the monthly mean data of global network of cosmic ray neutron monitor stations having different cut-off rigidity. In September, 2014, new results with almost twice as much data were presented of cosmic rays, in talk at CERN and published in Physical Review Letters (Schirber & Michael 2014).

Data Collection sources - For this work we have taken yearly average value of sunspot number from National Oceanic Atmospheric Administration (NOAA), their website is www.ngdc.noaa.gov and cosmic ray by Moscow neutron monitor having magnetic cut-off rigidity ($R_c = 2.42$ GeV) which is located on Earth at latitude 55.47N as well as longitude 37.32E similarly, Oulu neutron monitor have $R_c = 08$ GeV and location on earth at latitude 65.05N and longitude 25.47E.

Result and Discussion - The Neutron monitors are more sensitive to cosmic ray in the energy range 0.5-20 GeV, which coincides with maximum energy response for effective solar modulation. Though, the anti-correlation is found to vary during the different phases of solar cycles (Nagashima & Morishita 1979; Dorman *et al.* 2001). The sunspot number is the main characteristics of solar cycle variation. Sunspots are temporarily phenomenon on the photosphere of the sun and it's appearing visibly as dark spot compared to surrounding regions (Schwabe 1843). Solar cycle is also called solar activity magnetic cycle. The sunspots are most obvious features of photosphere and firstly observed by Galileo. The sunspots are composed of hot gases but they appear dark because the temperatures in the spots are a few thousand degree Kelvin cooler than that of surrounding photosphere. The Sunspot number varies from year to year with a certain regularity of increase and decrease. Cosmic ray flux comes from outward expansion of solar ejecta into interplanetary space. The



cosmic ray flux inters solar system anti-correlated with the solar activity. This anti-correlation is detected at the Earth surface by measurement. The cosmic ray flux varies with solar cycle during the period of solar cycle 22 to 24 (Anath 1975). The unit of cosmic ray intensity is (particle/- ster. sec. MeV). When cosmic rays flux inter solar system these flux modulated by the solar ejecta (Nagashima & morishita 1979; Mavoromichalaki & Petropoulos 1984; Nymmik & Suslov 1995; Storini *et al.* 1995). The cosmic rays are deflected by the magnetic field in interstellar space, they are also affected by the interplanetary magnetic field embedded in solar wind and therefore have difficulty reaching the inner solar system. As solar activity varies, over the 11 - year solar cycle the intensity of cosmic rays at Earth also varies, in anti-correlation with the sunspot numbers.




Fig. (1) Has shown the long - term change of state of the sun during period of solar cycle 22 to 24. Its modulate the cosmic rays during the same period of time. We found that cosmic ray intensity during solar cycle 24 have been increased than solar cycles 22 and 23.




Fig. 2. Has show the linear correlation of yearly average vale of sunspot number with cosmic ray intensity (Moscow and Qulu) during period of solar cycle 22 to 24 from year 1992 to 2022.

Fig. (2) Shown the linear correlation of sunspot number with cosmic ray intensity of Moscow neutron monitor much higher than cosmic ray intensity of Oulu neutron monitor. It means the both two neutron monitor have situated at different latitude and cut - off rigidity.

Conclusions - The graphical result obtained since a long period of time is showing that the activity of Sun has decreased during period of solar cycle 22 to 24. Therefore the solar phenomena i.e. solar temperature, proton flux, solar wind velocity, solar magnetic field and solar plasma should also decrease during the period of solar cycle 22 to 24 so as its consequence the disturbance in space weather should also decrease which may cause irregularity in Earth atmosphere during this period.

References

1. Ahluwalia, H.S. 1998, J. Geophys. Res., 103,12103.
2. Ahluwalia, H.S., Wilson, M.D. 1996, J. Geophys. Res.. 101 (A3), 4879.
3. Anath, A.G. 1975, The time variation of cosmic ray intensity, Ph.D. Thesis, Gujrat university, Ahmedabad (India).
4. Dorman, I.V., Dorman, L.I. 1967, J. Geophys. Res., 72 , 1513.



5. Dorman, L.I., Dorman, I.V., Lucci, N., Parisi, Villoresi, G. 2001, Adv. Space Res., 27, 589.
6. Gupta, M., Mishra, V.K., Mishra, A.P. 2005, Proc. 29th Intern. Cosmic Ray Conf., Pune, (India), 2,147.
7. Gupta, M., Mishra, V.K., Mishra, A.P. 2006, Indian J. Radio and Space Phys., 35,167.
8. Mavoromichalaki, H., Petropoulos, B. 1984, Asrophys. Space Sci., 106, 61.
9. Nagashima, K., Morishita, I. 1979, Proc. 16th Intern. Cosmic Ray Conf. Japan, 3, 325.
10. Nymmik, R.A., Suslov, A.A. 1995, Adv. Space Res., 16,217.
11. Parker, E.N. 1964, Planetary Space Sci. 12, 735.
12. Pomerantz, M.A. Duggal. S.P. 1971, J. Geophys. Res., 12,75.
13. Rao, U.R. 1972, Space Sci. Rev., 12,719.
14. Schwabe, S.H. 1843, Astronomical Nachrichten, 21, 233-236.
15. Schirber, Michael, “Synopsis: More dark matter hints from cosmic rays.” American physical society retrieved 21 sep 2014
16. Storini, M., Borello Filisetti, On the Mussino, N. Parisi, M., Sykora, J. 1995, Solar phys., 157, 375.
17. Usoskin, I.G., Kananen, H., Mursula, K., Tanskanen, P. 1998, J. Geophys. Res., 103, 9567.
18. Webber, W.R., Lockwood, J.A. 1988, J.Geophys. Res., 93,8735.

