

Effects of Cognitive Sensory Motor Training Versus Repetitive Facilitation Exercises of Upper Limb in Hemiparetic Patients

Ashish Patel, Department of computer science, R D engineering College, Ghaziabad
Corresponding author-patel.ashish@gmail.com

ABSTRACT

Stroke [CVA] is the sudden loss of neurological function caused by an interruption of blood flow to the brain. Large numbers of people who survive a stroke are left with permanent impairment of arm and hand function, even after completion of conventional rehabilitation programs. The standard neuro physiological facilitation technique used for hemiplegic upper limb have not been confirmed to promote functional recovery of hemiplegic limb. This promotes that more research needs to be conducted for same.

Cognitive Sensory Motor Training Therapy & The repetitive facilitation exercises (RFEs) Both techniques will promote functional recovery of hemiparetic upper limb and hand by improving joint perception and realization of movement. Hence, this study aims to compare the effectiveness of cognitive sensory motor training versus repetitive facilitation exercises on quality of movement of upper limb, functional activity and Range of motion of upper limb in hemiparetic patients.

A blinded randomized clinical trial was conducted 30 patients were divided into 2 groups (GROUP A and GROUP B)- those who performed cognitive Sensory Motor exercises (GROUP A- experimental group) and those who performed repetitive facilitation exercises (GROUP B-control group) Data for measures quality of movement performance of the hemiparetic arm and hand on MESUPES scale, Barthel Index (BI) measures the extent to which somebody can function independently and has mobility in their &, goniometer measuring the joint ranges in each plane of the joint was collected on day 1 (pretreatment session), and on 190 day after the experiment.

This study produced a statistically significant increase in overall on quality of movement, functional activity and range of motion in both the group of upper limb in hemiplegic patients. This research also provides evidence that training exercise may be a valuable and important tool in clinical practice and is consistent with the current use by clinical physiotherapist in the treatment of upper limb in hemiplegic patients.

Keywords: Stroke, Barthel Index, MESUPES scale

INTRODUCTION

Stroke (cerebrovascular accident [CVA]) is the sudden loss of neurological function caused by an interruption of the blood flow to the brain. Ischemic stroke is the most common type, affecting about 80% of individuals with stroke, and results when a clot blocks or impairs blood flow, depriving the brain of essential oxygen and nutrients. Hemorrhagic stroke occurs when blood vessels rupture, causing leakage of blood in or around the brain. Clinically, a variety of focal deficits are possible, including changes in the level of consciousness and impairments of sensory, motor, cognitive, perceptual, and language functions. To be classified as stroke, neurological deficits must persist for at least 24 hours. Motor deficits are characterized by paralysis (hemiplegia) or weakness (hemiparesis), typically on the side of the body opposite the side of the lesion. The term hemiplegia is often used generically to refer to the wide variety of motor problems that result from stroke. The location and extent of brain injury, the amount of collateral blood flow, and early acute care management determine the severity of neurological deficits in an individual patient. Impairments may resolve spontaneously as brain swelling subsides (reversible ischemic neurological deficit), generally within 3 weeks. Residual neurological impairments are those that persist longer than 3 weeks and may lead to lasting disability. Strokes are classified by etiological categories (thrombosis, embolus, or hemorrhage), specific vascular territory (anterior cerebral artery syndrome, middle cerebral artery syndrome, and so forth), and management categories (transient ischemic attack, minor stroke, major stroke, deteriorating stroke, young stroke).

PREVALENCE AND INCIDENCE OF STROKE

Stroke is the fourth leading cause of death and the leading cause of long-term disability among adults in the United States. An estimated 7,000,000 Americans older than 20 years of

age have experienced a stroke. Each year approximately 795,000 individuals experience a stroke; approximately 610,000 are first attacks and 185,000 are recurrent strokes. Women have a lower age-adjusted stroke incidence than men. However, this is reversed in older ages; women over 65 years of age have an elevated risk compared to men. Compared to whites, African Americans have twice the risk of first- ever stroke; rates are also higher in Mexican Americans, American Indians, and Alaska Natives. The incidence of stroke increases dramatically with age, doubling in the decade after 65 years of age. Twenty-eight percent of strokes occur in individuals younger than 65 years of age. Between 5% and 14% of persons who survive an initial stroke will experience another one within 1 year; within 5 years stroke will recur in 24% of women and 42% of men. Current data reveal that stroke incidence has been declining in recent years in a largely white adult cohort.¹ The incidence of stroke deaths is greater than 143,000 annually, and strokes account for 1 of every 18 deaths in the United States. The type of stroke is significant in determining survival. Of patients with stroke, hemorrhagic stroke accounts for the largest number of deaths, with mortality rates of 37% to 38% at 1 month, whereas ischemic strokes have a mortality rate of only 8% to 12% at 1 month. Survival rates are dramatically lessened by increased age, hypertension, heart disease, and diabetes. Loss of consciousness at stroke onset, lesion size, persistent severe hemiplegia, multiple neurological deficits, and history of previous stroke are also important predictors of mortality.

Stroke is the most common cause of chronic disability. Of survivors, majority will experience difficulty with activities of daily living (ADLs), ambulation, speech, motor disturbance, sensory disturbance, perceptual disturbance, language disturbance, cognitive disorder, and urinary incontinence depending on the area of the brain lesion. Hemiplegia is commonly associated with a decrease in balance ability. 50% to 65% of stroke patients are left with functional impairments. Most patients are still significantly disabled beyond 6 months after stroke, and do not return to social activities within the community. Large numbers of people who survive a stroke are left with permanent impairment of arm and hand function, even after completion of conventional rehabilitation programs. It has been reported that only 5–20% of patients regain full arm and hand function with a number of prospective cohort studies suggesting that 33–66% of stroke patients with a paretic arm do not show any recovery of upper limb function six months after stroke.

The signs and symptoms of stroke are as follows

- Hemiparesis and weakness of facial muscles
- Numbness
- Altered sensation
- Initial flaccidity of muscles (decreased tone of muscles) which is later replaced by spasticity (increase in tone of muscles), exaggerated reflexes and development of synergies. Majority of the cases of stroke represent unilateral weakness that is weakness on one side of the body. Due to inter-crossing of the fibres of the brain the symptoms usually appear on the opposite side of the area of brain being affected. The human brain is divided into forebrain, midbrain and hindbrain. Forebrain consists of the cerebrum and hindbrain comprises of medulla oblongata, pons verolli and cerebellum. Depending on the severity and extent of damage occurred in the brain different signs and symptoms are seen which are explained as follows-
- Altered or disturbed sense of smell, taste, hearing or vision.
- Disturbed visual fields
- Weakness of ocular muscles, characterised by drooping of eyelids
- Decreased reflexes primarily the gag reflex, swallow and reactivity of pupil towards light
- Altered sensation on extremities and weakness of the facial and axial musculature.
- Difficulty in balancing and development of nystagmus
- Difficulty in articulation
- Drooling of saliva

Large numbers of people who survive a stroke are left with permanent impairment of arm and

hand function, even after completion of conventional rehabilitation programs. It has been reported that only 5–20% of patients regain full arm and hand function with a number of prospective cohort studies suggesting that 33–66% of stroke patients with a paretic arm do not show any recovery of upper limb function six months after stroke.

TREATMENT OF HEMIPLEGIC UPPER LIMB

Understanding upper limb impairment after stroke is essential to planning therapeutic efforts to restore function. However, determining which upper limb impairment to treat and how is complex for two reasons:-

1) The impairments are not static, i.e., as motor recovery proceeds, the type and nature of the impairments may change; therefore, the treatment needs to evolve to target the impairment contributing to dysfunction at a given point in time.

2) Multiple impairments may be present simultaneously, i.e., a patient may present with weakness of the arm and hand immediately after a stroke, which may not have resolved when spasticity sets in a few weeks or months later; hence there may be a layering of impairments over time making it difficult to decide what to treat first.

The most useful way to understand how impairments contribute to upper limb dysfunction may be to examine them from the perspective of their functional consequences.

There are three main functional consequences of impairments on upper limb function are: (1) learned nonuse, (2) learned bad-use, and (3) forgetting as determined by behavioral analysis of tasks. The impairments that contribute to each of these functional limitations are described. Cognitive Sensory Motor Training Therapy is a unique comprehensive rehabilitation programme incorporating systematic coaching and retraining of sensory guided motor control.

First proposed by Professor Carlo Perfetti, this rehabilitation programme is now known as Perfetti's Method. Perfetti's Cognitive Sensory Motor

Training Therapy is that it focuses on sensory retraining, with particular emphasis on joint position perception.

The repetitive facilitation exercises (RFEs) using novel facilitation methods for the upper limb and fingers, give sufficient physical stimulation, such as by the stretch reflex or skin-muscle reflex that is elicited immediately before or at the same time as when the patient makes an effort to move his hemiplegic hand or finger, in order to elevate the level of excitation of the corresponding injured descending motor tracts and it allows the patient to initiate movements of the hemiplegic hand or finger in response to his intention.

Limitations in arm and hand function are a major problem after stroke and cause difficulties in patients' daily lives. Recent research has demonstrated that the adult central nervous system retains a much higher capacity for plasticity and reorganization than earlier believed, therefore, an important goal of stroke rehabilitation is to substantially increase the functional use of the affected arm while minimizing compensatory strategies and avoiding learned disuse.

METHODOLOGY AIM

To compare the effectiveness of cognitive sensory motor training versus repetitive facilitation exercises on quality of movement of upper limb, functional activity and Range of motion of upper limb in hemiparetic patients.

OBJECTIVE

To identify the effect of Cognitive Sensory Motor Training in individuals with Hemiparetic patients. To identify the effect of Repetitive Facilitation Exercise in individuals with Hemiparetic patients. To investigate the effect of cognitive sensory motor training versus repetitive facilitation exercises on quality of movement of upper limb, functional activity and Range of motion of upper limb in hemiparetic patients.

SAMPLING

Study Design

A comparative study **Sample Method** Experimental

Sample size

The sample size will consist of 30 hemiparesis patients with the equal size of 15 patients in each of the two groups.

Samplesetting

The study will be originated at career hospital bhopal.

Study Duration


3 month

Inclusion criteria

- Gender: Both male and female
- Age: 25-65 years
- Patient with stroke confirmed by MRI no previous history of stroke
- Mini Mental State Examination Score should be >21 OUT OF 30
- Brunnstrom Recovery Stage Score should be ≥ 4
- Stroke within less than 4 weeks before the study.
- Confirmed by clinical examination and magnetic resonance imaging (MRI).
- Ability to follow simple direction of commands.

Exclusion criteria

- Patient with severe sensory disturbance, pain and contracture.
- Patient with hemineglect pre-existing upper extremity impairment.
- Patient without motor deficits.
- Presence of any other musculoskeletal condition. e.g. (Frozen shoulder, any recent fractures of upper limb).
- Any accompanying diseases or disorders, other than stroke, that could interfere with upper extremity training.
- Uncontrolled health conditions for which exercise was contraindicated.
- Cerebellar lesion.



VARIABLES

Dependent variables

1. MESUPES scale
2. Modified Barthel Index (MBI)
3. Goniometer

Independent variables

1. Arm Function
2. Visual Analogue Scale
3. Muscle power

INSTRUMENTATION AND FUNCTIONAL SCALES USED

1. MESUPES scale
2. Barthel index
3. Goniometer set
4. Chair
5. Table
6. Dice

PROCEDURE

Exercise protocols

A blinded randomized clinical trial was conducted 30 patients were divided into 2 groups (GROUP A and GROUP B)- those who performed cognitive Sensory Motor exercises (GROUP A- experimental group) and those who performed repetitive facilitation exercises (GROUP B- controlled group) Data for measures quality of movement performance of the hemiparetic arm and hand on MESUPES scale, Modified Barthel Index (MBI) measures the extent to which somebody can function independently and has mobility in their & Goniometer measuring the joint ranges in each plane of the joint was collected on day 1 (pretreatment session), and on 90 day after the experiment.

Both groups underwent their respective interventions for 30 min each in the morning and 30 min in the afternoon to minimize the physical fatigue.

Measurements MESUPES scale

This approach has been used in the development of the new assessment tool, the Motor

Evaluation Scale for Upper Extremity in Stroke Patients (MESUPES).A 17-items into two sub scales ; MESUPES- Arm function; 8items(score 0-5) MESUPES-Hand function;9items(score0-2) objective evaluation scale designed to assess quality of movement of armand hand function after stoke.

ModifiedBarthelIndex (MBI)

The MBI, which consists of 10 items describing activities of daily living (ADL) and mobility, wasscored to measure the degree of assistance required by an individual and was used to assess ADL in patients with stroke . Each item is rated 5-Likert scale, with weights added according to the item. The higher the total score, the more independent on performing ADLs.

Goniometer

The rangeofmotionisthemasurementofmovementaroundaspecificjointrangeofmotion,A universal goniometer hasthreeparts.Abody-Itisdesignedlikeaprotractorandmayformafullora half-circle. It has a scale for the measurement of the angle. The scale can extend either from 0 to 180 degrees or 180 to 0 degrees for half circle models or 0 to 360 degrees on full circle models. The moving arm is the arm of the goniometer, which aligns with the mobile part of the joint measured.

Intervention

GroupA(experimentalGroup)CognitiveExerciseTherapy

PRE

Day1

20repsx2set

1. Shoulder joint recognition training by motor imagery
2. Shoulder and elbow joint recognition training using a circular track plate.
3. Training on awareness of elbow and wrist joint angles using a Bogen. Training on pressure awareness of the elbow and wrist using a sponge.
4. Fingertactile recognition training using a tactile plate.

GroupA(experimentalGroup)CognitiveExerciseTherapy

POST

Day90 20repsx2set

1. Shoulder joint recognition training by motor imagery
2. Shoulder and elbow joint recognition training using a circular track plate
3. Training on awareness of elbow and wrist joint angles using a Bogen.
4. Training on pressure awareness of the elbow and wrist using a sponge.
5. Fingertactile recognition training using a tactile plate.

GROUPB(controlgroup)RepetitiveFacilitationExercises Each RFE session includes eight specific exercise patterns. PRE DAY1

20repsx2set

1. Shoulder flexion with 90 elbow flexion.
2. Shoulder horizontal extension/flexion with elbow flexion.
3. Shoulder flexion/adduction/external rotation with flexion of the elbow and forearm supination with wrist flexion, finger flexion followed by shoulder extension/abduction/internal rotation while extending the elbow and pronating the forearm accompanied by wrist dorsiflexion and finger extension in the supine position.
4. Shoulder flexion/abduction/external rotation with elbow extension accompanied by wrist extension and finger extension (modified PNF).
5. Forearm supination/pronation with 90 elbow flexion in the sitting position When the therapist will give commands,, Turn your hand (palm) upward", the patient attempts to perform forearm supination and then ask to,, Turn your hand (palm) down-ward", the patient attempts to perform forearm pronation.
1. Wrist extension and forearm pronation with extension of the fingers in the supine position.
2. Finger extension with wrist flexion in the supine position.
3. Finger extension/flexion with wrist flexion in the sitting position.

GROUPB(controlgroup)RepetitiveFacilitationExercises

POSTDAY90

1. shoulderflexionwiththeelbowbentat90°inthesupineposition
2. shoulder horizontal extension/flexion in the supine position with the elbow ranging in flexion from about 70° to 110°
3. shoulder flexion/adduction/external rotation with flexion of the elbow and forearm supination accompanied by wrist flexion, finger flexion, and shoulder extension/abduction/internal rotation while extending the elbow and pronating the forearm accompanied by wrist dorsiflexion and finger extension in the supine position
4. shoulder flexion/abduction/external rotation with elbow extension accompanied by wrist dorsiflexion and finger extension
5. forearmsupination/pronationwith90°elbowflexioninthesittingposition
6. wristdorsiflexionandforearmpronationwithextensionofthefingersinthesupineposition
7. fingerextensionwithwristflexioninthesupineposition
8. fingerextension/flexionwithwristflexioninthesittingposition

Conventional therapy

Control activities consisted of self-range of motion (SROM) stretches and active range of motion (AROM) strengthening exercises throughout the hemiparetic upper extremity.

During SROM stretches, participants clasped the hands or arms together and used the strength of the less-affected arm to move the affected arm through the available ROM at each joint.

During AROM exercises, the hemiparetic arm was supported against gravity by a tabletop, and a towel was placed under the arm.

DISCUSSION

The present study was undertaken with the intention to see the effectiveness of Cognitive Sensory Motor Training Versus Repetitive Facilitation Exercise for Quality of movement, ADL and ROM in Subjects with Hemiparesis (age 25 to 65 years) using MESUPES and BI scales. The MESUPES is a tool used to check quality of movement in hemiparetic upper limb, while the BI provides guidelines for determining Daily living activities and functional levels and treatment. A total of 30 subjects both males and females aged 25 to 65 years with stroke (Hemiparesis) were included with 15 participants in each group out of which a total of 15 subjects in group A and 15 subjects in group B completed the 12 weeks of program.

In this study we found that Cognitive Sensory Motor Training is effective in improving Quality of movement of upper limb, functional activity and ROM ($p \leq 0.0001$) in Hemiparetic patients. Similarly, Repetitive Facilitation Exercise for is also effective but not more than cognitive sensory motor exercise in improving Quality of movement of upper limb, functional activity and ROM ($p < 0.0001$) in Hemiparetic patients.

The results of the present study are in agreement with the study conducted by Ratanapat Chanubolet al, (2012) studied the effectiveness of Cognitive Sensory Motor Training Therapy (Perfetti's method) vis-à-vis conventional occupational therapy in the recovery of arm function after acute stroke by Prospective randomized controlled trial in rehabilitation centers in Bangkok, Thailand.

The better improvement in cognitive sensory motor training is because it focuses on sensory retraining, with particular emphasis on joint position perception, incorporating systematic coaching and retraining of sensory guided motor control. And The repetitive facilitation exercises (RFEs) using novel facilitation methods for the upper limb and fingers, give sufficient physical stimulation, such as by the stretch reflex or skin-muscle reflex that is elicited immediately before or at the same time as when the patient makes an effort to move his hemiplegic hand or finger, in order to elevate the level of excitation of the corresponding injured descending motor tracts and it allows the patient to initiate movements of the hemiplegic hand or finger in response to his intention.

CONCLUSION

The common trend of the treatment of functional activity in hemiparetic upper limb. Exercises for hemiparetic upper limb concentrate on increasing range of motion with the assumption that functional improvement will follow. In this study, the cognitive sensory motor group demonstrated an increase in quality of movement level by (75%) when compared to the repetitive facilitation exercise group (50%), which thus could account for the functional ability

differences seen between groups. In addition to this the observed group effect for functional activity and ROM in favor of the cognitive sensory motor exercise may have been key in improving functional stability. It is necessary to recognize if deficits at the impairment level are causative in limiting activities, so that if strength is an issue, dealing with the impairment at a more functional level may be more effective in the long term. Effectiveness of strengthening exercises can be maximized by introducing flexibility, coordination, balance, and mobility, which may transfer to an overall improvement in function. Repetition and task practice not only improves strength but reduces activity limitations associated with the impairment of decreased muscle strength. Ultimately, the inability to participate in activities at a social level has an impact on the quality of life in individuals with hemiplegic patients. Addressing the impairments and activity limitations associated with this disease in middle-aged individuals may delay and/or prevent the disabilities encountered in the elderly. One can also suggest that the repetition of the star exercise contributed to proprioceptive acuity and increased balance and stability in the cognitive sensory motor group as it involves balancing on upper limb while reaching out with the other upper limb to touch all points of an outlined star. The data from this study support that cognitive sensory motor exercise is a better option in improving quality of movement of upper limb, functional activity and ROM in this 25- 65 year old population with Hemiparetic patients.

REFERENCES

1. Susan O'sullivan B. Physical rehabilitation, 6th edition, chapter 15, 645
2. Pandian JD, Sudhan P. Stroke epidemiology and stroke care services in India. Journal of Stroke 2013
3. Chanubol R, Wongphaet P, Chavanich N, Werner C, Hesse S, Bardeleben A, et al. A randomized controlled trial of Cognitive Sensory Motor Training Therapy on the recovery of arm function in acute stroke patients. Clinical rehabilitation 2012
4. Kawahira K, Shimodzono M, Etoh S, Kamada K, Noma T, Tanaka
5. N. Effects of intensive repetition of a new facilitation technique on motor functional recovery of the hemiplegic upper limb and hand. Brain Injury 2010
6. Johansson GM, Häger CK. Measurement properties of the motor evaluation scale for upper extremity in stroke patients (MESUPES). Disability and rehabilitation 2012
7. Pendleton HM, Schultz-Krohn W. Pedretti's Occupational Therapy-E- Book: Practice Skills for Physical Dysfunction. Elsevier Health Sciences 2017,
8. Bour A, Rasquin S, Boreas A, Limburg M, Verhey F. How predictive is the MMSE for cognitive performance after stroke?. Journal of neurology 2010
9. Housman SJ, Scott KM, Reinkensmeyer DJ. A randomized controlled trial of gravity-supported, computer-enhanced arm exercise for individuals with severe hemiparesis. Neurorehabilitation and neural repair 2009
10. Hsueh IP, Lee MM, Hsieh CL. Psychometric characteristics of the Barthel activities of daily living index in stroke patients. Journal of the Formosan Medical Association. 2001
11. Hiengkaew V, Vittayasoontorn P, Meenathanin P, Kaewtong A. Joint Range of Motion in Flaccid Hemiplegia. Hong Kong Physiotherapy Journal 2003
12. Rabadi MH. Review of the randomized clinical stroke rehabilitation trials in 2009. Medical science monitor: international medical journal of experimental and clinical research 2011
13. Wongphaet P, Butrach W, Sangkrai SI, Jitpraphai C. Improved function of hemiplegic upper extremity after cognitive sensory motor training therapy in chronic stroke patients: preliminary report of a case series. Journal of the Medical Association of Thailand=Chotmaihetthangphaet 2003
14. Feys H, De Weerd W, Verbeke G, Steck GC, Capiau C, Kiekens C, et al. Early and repetitive stimulation of the arm can substantially improve the long-term outcome after stroke: A 5-year follow-up study of a randomized trial. Stroke 2000.
15. Langhammer B, Stanghelle JK. Bobath or motor relearning programme? A comparison of two different approaches of physiotherapy in stroke rehabilitation: a randomized controlled study. Clinical rehabilitation. 2000

16. Duncan PW, Goldstein LB, Horner RD, Landsman PB, Samsa GP, Matchar DB. Similar motor recovery of upper and lower extremities after stroke. *Stroke* 1994;25(6):1181-8.
17. Nakayama H, Jørgensen HS, Raaschou HO, Olsen TS. Recovery of upper extremity function in stroke patients: the Copenhagen Stroke Study. *Archives of physical medicine and rehabilitation*. 1994
18. Sunderland A, Fletcher D, Bradley L, Tinson D, Hewer RL, Wade DT. Enhanced physical therapy for arm function after stroke: a one year follow up study. *Journal of Neurology, Neurosurgery & Psychiatry*. 1994
19. Lopez A.D., Mathers C.D., Ezzati M., Jamison D.T., Murray C.J. Global and regional burden of disease and risk factors, 2001
20. Aman J.E., Elangovan N., Yeh I.L., Konczak J. The effectiveness of proprioceptive training for improving motor function: A systematic review. *Front. Hum. Neurosci.* 2015
21. Lee S., Bae S., Jeon D., Kim K.Y. The effects of cognitive exercise therapy on chronic stroke patients' upper limb functions, activities of daily living and quality of life. *J. Phys. Ther. Sci.* 2015

